

SOFT CONTROL OF SWARMS - ANALYTICAL APPROACH

Guillaume Sartoretti*¹ and Max-Olivier Hongler¹

¹*STI/IMT/LPM, École Polytechnique Fédérale de Lausanne (EPFL), CH-1015, Lausanne, Switzerland*
{guillaume.sartoretti,max.hongler}@epfl.ch

Keywords: Rank-Based Brownian Motions - Heterogeneous autonomous agents - Soft-control of swarms - Exact analytic solvable models - Super-diffusive skill agent - Flocking

Abstract: We analytically study the collective dynamics of mutually interacting heterogeneous agents evolving in a random environment. Our formal framework consists of a collection of N scalar drifted Brownian motions (BM) diffusing on \mathbb{R} . The mutual interactions are introduced via a ranked-based, real-time mechanism always endowing the laggard (i.e the agent with the leftmost position) with an extra positive drift. The extra drift generates a net tendency for any agents not to remain the laggard of the society. For well chosen individual and extra laggard's drifts, the agents organize with time to flock towards a tight and stable traveling spatial pattern. For a population of $(N - 1)$ identical agents and an atypical fellow (called hereafter the skill), we are able to analytically discuss the dynamics. In particular we exhibit how a single turbulent skill, stylized here by a ballistic diffusion process, can destroy the cohesion of a swarm. Conversely, we also analytically show how a single skill is able to safely pilot a whole swarm to avoid an obstacle, via interactions with its fellows. A series of simulations experiments comfort our analytic findings.

1 INTRODUCTION

The capability of a collection of interacting stochastic agents to exhibit an emergent collective behavior (i.e *flocking behavior*) even in random environments stimulates a strong research activity devoted to both experimental and theoretical modeling approaches. For suitable range of mutual interactions, flocking (phase) transitions are observed, namely the self-organized capability to create finite and persistent spatio-temporal patterns (Banner A. D. and Karatzas, 2005; Yates C. A. and Sumpter, 2009; Chatterjee and Pal, 2010; Dosseti, 2012; Pal and Pitman, 2008; Ichiba T. and Fernhold, 2011; Bialek W. and Walczak, 2012).

Agents societies can be composed of either dynamically homogeneous or heterogeneous individuals requiring for each case drastically different approaches. For large and **homogeneous** population of agents, the classical statistical mechanics concepts and in particular the **mean-field description** (MF) directly offers an appropriate tool to analytically discuss the global dynamics. In the MF description, one basically assumes that the behavior of the global society can be characterized by the dynamics of a single representa-

tive agent which feels its fellows' interactions via an effective external field. In most circumstances however, homogeneity fails, and therefore a growing attention is now paid to **heterogeneous** populations. Heterogeneity occurs typically when one or several masked agents, that we shall from now on refer to as **skills**, exhibit leaders or troublemakers behaviors. These skills are not recognized by the regular agents which see them as ordinary fellows, leaving the interaction rules between them unchanged. The presence of skills can strongly alter the ultimate evolution of the whole society and it is the central goal here to analytically study this problematic.

Our approach involves assemblies of interacting stochastic agents in which a single skill exhibits a different individual behavior. All agents, including the skill, interact with their fellows with fixed given rules. The paradigmatic vision of this situations has been currently explored in ethology where one fake individual is introduced among schools of fishes, cockroaches, newborn chicken, etc. The skill is able to ultimately pilot the whole population (Gribovskiy A. and Mondada, 2010; Faria J. J. and Krause, 2010; Vaughan R. and Cameron, 2000). This basic mechanism also referred as the **soft control** of a population (Han and Wang, 2010; Wang X. and Han, 2011; Wang Y. and Liu, 2009; Wang and Guo, 2008) is the

*Supported by the Swiss National Funds for Scientific Research

core of our present paper. As heterogeneous societies preclude MF approaches, rather few analytical results are yet available. Recent mathematical models of heterogeneous **rank-based interacting Brownian motions** (RBM), introduced in finance (Banner A. D. and Karatzas, 2005; Chatterjee and Pal, 2010; Pal and Pitman, 2008; Ichiba T. and Fernhold, 2011), will be used in the sequel to analytically approach the soft control problematic. We shall **analytically show** how a **troublemaker can break the cohesion** of an initially tight swarm and **how a single agent can efficiently incite its fellows to achieve a preassigned task**.

Our paper is organized as follows: we first recall in section 2 relevant properties of RBM, also known in economy as the **Atlas models**. In section 3, we use the RBM's framework to show how a troublemaker is able to smash an initially tight swarm. Our individual dynamics are one-dimensional diffusion processes with piecewise constant drifts except for the skill, which will be assumed to be driven by a **ballistic process with quadratic variance in time**. The ballistic noise is itself generated by a simple non-Gaussian diffusive stochastic process with nonlinear drift. In section 4, we address the dual soft control problem in which a skill is used to steer the whole collection of RBM's towards slots that are drilled through an obstacle board.

2 RANK-BASED BROWNIAN MOTIONS - ATLAS MODEL

Our approach makes extensive use of recent results (Ichiba T. and Fernhold, 2011), that we now briefly summarize. Let us consider a collection of N interacting agents diffusing according to the class of processes:

$$dY_i(t) = \left(\sum_{k=1}^N g_k 1_{Q_k(i)} \{Y(t)\} + \gamma_i + \gamma \right) dt + \sigma_i dW_i(t),$$

$$Y_i(0) = y_i, \quad t \in \mathbb{R}^+, \quad (1)$$

where $Y(t) = (Y_1, Y_2, \dots, Y_N) \in \mathbb{R}^N$ and $dW_i(t)$ are N independent standard White Gaussian Noise processes (WGN) processes. The indicator function 1_{Q_k} in Eq.(1) effectively generates mutual interactions. The effective, time-dependent drift component g_k entering into the drifts $(g_k + \gamma_i + \gamma)$ of the N Brownian motions on \mathbb{R} , is **rank-based**. Namely, it is instantaneously adjusted according to the position (i.e. the rank) occupied by each agent with respect to the remaining $(N - 1)$ fellows. The constant drift

components γ_i are name-based, i.e. they are definitely assigned to each individual agent (γ_i is time-independent). Finally, a constant drift component γ can be added, which is common to all agents. Accordingly Eq.(1) describes a collection of N diffusion processes having piecewise deterministic drifts. The somehow simpler situation obtained when $\gamma_i \equiv 0 \forall i$ has been thoroughly studied (Pal and Pitman, 2008). In the sequel and without loss of generality, we systematically choose γ to be the (average) barycentric speed of the swarm. This is achieved, provided one has:

$$\sum_{k=1}^N [g_k + \gamma_k] = 0. \quad (2)$$

For future use, we introduce the following notations and definitions:

1. We write

$$\bar{Y}(t) = \frac{1}{N} \sum_{k=1}^N Y_k(t), \quad \bar{Y} \in \mathbb{R}, \quad (3)$$

for the barycenter position. The set Σ_N stands for the set of all the permutations of $\{1, 2, \dots, N\}$. It is proven (Ichiba T. and Fernhold, 2011) that, for almost every initial conditions and when the set of constraints

$$\sum_{k=1}^l [g_k + \gamma_{p(l)}] < 0 \quad (4)$$

are fulfilled for all possible permutations $p = (p(1), \dots, p(N)) \in \Sigma_N$, then the N deviations processes:

$$\tilde{Y}_i(t) = [Y_i(t) - \bar{Y}(t)] \quad (5)$$

converge to stationary probability measures. Note that Eq.(4) yields therefore a set of $(N - 1) \times |\Sigma_N| = N! (N - 1)$ constraints to be verified.

2. When all constraints in Eq.(4) are fulfilled, the dynamics given by Eq.(1) then converges to a **tight swarm** described by a **stationary multi-variate process** characterizing the $(N - 1)$ -gap processes

$$\Xi_i(t) = [Y_{i+1}(t) - Y_i(t)]. \quad (6)$$

The associated probability density $\psi(z)$, $z \in \mathbb{R}_+^{N-1}$ can be written as a sum-of-product-of-exponential form:

$$\psi(z) = \left(\sum_{p \in \Sigma_N} \prod_{k=1}^{N-1} \lambda_{p,k}^{-1} \right)^{-1} \sum_{p \in \Sigma_N} \exp(-\langle \lambda_p, z \rangle) \quad (7)$$

with the parameters explicitly given by

$$\lambda_p = (\lambda_{p,k})_{k=1}^{N-1} \quad \lambda_{p,k} = \frac{-4 \sum_{l=1}^k (g_l + \gamma_{p(l)})}{\sigma_k^2 + \sigma_{k+1}^2}. \quad (8)$$

Remark. It is worth observing that the diffusion coefficients σ_i do not enter into the set of tightness constraints given in Eq.(4). However, the σ_i 's do enter into the parameters Eq.(7) characterizing the stationary probability measure.

3 HOW DOES A SUPER-DIFFUSIVE FELLOW SMASH TIGHTNESS

We first investigate how a single "turbulent" fellow can destroy the cohesion of a tight swarm. To this aim, we use the RBM formalism introduced in section (2). We consider a configuration involving $(N - 1)$ identical mutually interacting RBM's, referred from now on as the regular agents, interacting with a single fellow, the **skill**, say agent number one, which is itself driven by a **ballistic diffusion process** (remember from Eq.(1) that the $(N - 1)$ regular agents are driven by independent WGN's). The ballistic process, to be introduced below, exhibits a variance $\propto t + \beta t^2$ with a **ballistic parameter** $\beta \geq 0$ a constant. For $\beta = 0$, the skill simply behaves as a regular fellow and therefore, in this $\beta = 0$ limit, our global dynamics reduces to a standard version of Eq.(1). We set the specific parameters: $g_N = (N - 1)g$ and $g_k = -g$ for $1 \leq k \leq N - 1$. In addition, we shall further assume that:

- i) $\sum_{i=1}^N \gamma_i = 0$,
- ii) $\max_{1 \leq i \leq N} \gamma_i < g$.

The couple of constraints i) and ii) imply that the Eq.(4) is fulfilled. Indeed, we directly verify that :

$$-g + \gamma_{p(1)} < 0, \quad -2g + \gamma_{p(1)} + \gamma_{p(2)} < 0, \quad \dots$$

$$-(N - 1)g + \sum_{k=1}^{N-1} \gamma_{p(k)} < 0. \quad (9)$$

Hence, for $\beta = 0$, our collection of agents behaves as a tight swarm enjoying a stationary joint probability law for the inter-distance between successive agents.

Let us now focus on $\beta > 0$. Now, the skill $Y_1(t)$ behaves as a **turbulent fellow** which interact with the remaining $(N - 1)$ regular fellows. Specifically the dynamics of $Y_1(t)$ is chosen to be described by:

$$dY_1(t) = \left(\sum_{k=1}^N g_k 1_{Q_k(i)} \{Y(t)\} + \gamma_1 + \gamma \right) dt + dZ(t),$$

$$Y_1(0) = y_1 \quad t \in \mathbb{R}^+, \quad (10)$$

where the process $dZ(t)$ is a **super-diffusive ballistic noise generator** introduced in (Hongler M. O. and Blanchard, 2006; Hongler M. O. and Rodriguez, 2012):

$$dZ(t) = \{\beta \tanh[\beta Z(t)]\} dt + dW(t), \quad Z(t=0) = 0. \quad (11)$$

The non-Gaussian Markov diffusion process $Z(t)$ given in Eq.(11) is fully characterized by its transition probability density $\mathcal{P}(z, t | 0)$ which simply reads:

$$\mathcal{P}(z, t | 0) = \frac{(\mathcal{P}_+ + \mathcal{P}_-)(z, t | 0)}{2} = \frac{e^{-\frac{\beta^2}{2}t} e^{\frac{(z \pm \beta t)^2}{2t}}}{\sqrt{2\pi t}}, \quad (12)$$

with average $\langle Z(t) \rangle = 0$ and variance $\langle Z^2(t) \rangle = t + \beta^2 t^2$.

The simple form given in Eq.(12) suggests the existence of an alternative representation for the ballistic noise $Z(t)$. Indeed, writing $BM_{\pm\beta}(t)$ for the Brownian motions with $\pm\beta$ constant drifts, we observe that the transition probability for the process $Z(t)$ can be rewritten as $Z(t) = \mathcal{B}BM_{\pm\beta}(t)$ where \mathcal{B} is a symmetric Bernoulli r.v., taking the values ± 1 with equal probability $1/2$. Hence, one realization of the $Z(t)$ consists first in choosing with probability $1/2$ one among the couple processes $BM_{\pm\beta}(t)$, and then follow the realization of the selected process (see example 2 in (Rogers and Pitman, 1981) and (Hongler M. O. and Blanchard, 2006; Hongler M. O. and Rodriguez, 2012)).

We now come back to the dynamics jointly involving a turbulent fellow given by Eq.(10) and $(N - 1)$ regular agents described by Eq.(1). We then view the $(N - 1)$ regular agents as being **infiltrated** by a **the skill** $Y_1(t)$.

Now we have to investigate the values of β enabling the swarm to remain tight (i.e. if a stationary probability measure for the intervals between successive agents exists). In view of the representation given in Eq.(12), for each realization of the noise source $Z(t)$, we effectively deal with a standard RBM model with a re-normalized name-based drift of $Y_1(t)$, namely $\gamma_1 \mapsto \gamma_1 \pm \beta$ depending on the outcome of \mathcal{B} . Accordingly, to infer on the tightness of the swarm, we simply have to separately examine Eq.(4) for the couple of outcomes $\pm\beta$.

Realization $+\beta$. Let us define $\gamma = \frac{\beta}{N}$, $\gamma_1 = \frac{N-1}{N}\beta$ and $\gamma_i = -\gamma = -\frac{\beta}{N}$ for $2 \leq i \leq N$. The constraints Eq.(6) are required for the swarm to be tight. This yields, for $l = 1$ and $p = (1, 2, 3, \dots, N)$, to the most critical constraint:

$$g_1 + \gamma_{p(1)} < 0 \quad \Rightarrow \quad -g + \frac{N-1}{N}\beta < 0$$

$$\Leftrightarrow \quad \beta < \frac{N}{N-1}g. \quad (13)$$

Provided that Eq.(13) holds, the swarm remains tight when the Z_t noise induces a $+\beta$ extra drift.

Realization $-\beta$. The same reasoning applied to the $-\beta$ case yields:

$$\gamma = -\frac{\beta}{N} \quad \gamma_1 = -\frac{N-1}{N}\beta \quad \gamma_i = \frac{\beta}{N} \quad (2 \leq i \leq N). \quad (14)$$

The critical constraint arises when $l = 1$ and $\gamma_p := \left(\frac{\beta}{N}, \dots, \frac{\beta}{N}, -\frac{N-1}{N}\beta\right)$ with:

$$-g + \frac{\beta}{N} < 0 \Leftrightarrow \beta < N \cdot g. \quad (15)$$

This implies that for $\beta < N \cdot g$, the swarm remains tight when the Z_t noise induces a $-\beta$ extra drift.

The previous considerations can be summarized by observing that when the constraints Eq.(13) are fulfilled, so are also those given in Eq.(15). This suggests to distinguish an alternative “**semi**”-tight regime which arises when the constraints Eq.(15) alone are fulfilled. In this “semi”-tight case, the swarm exhibits a tight configuration when the $-\beta$ realization is achieved and is not tight for the other alternative $+\beta$. This intrinsic asymmetry can be easily understood as our RBM dynamics only pushes the laggard towards the others. Hence, regarding the tightness, the skill is obviously less influential when the $-\beta$ extra drift is realized.

3.1 Spatial Dispersion of the Agents

The extra $\pm\beta$ -drifts due to the super-diffusive process driving the skill dynamics obviously affect the spatial dispersion of the swarm. As all noise sources (including the $dZ(t)$ process) have zero average, the barycenter of the whole population will remain unchanged by the presence of $dZ(t)$. However, due to the presence of the skill, an initial single-modal cluster of agents will, as time increases, be split into two separately evolving population subgroups. According to the $\pm\beta$ realization taken by $dZ(t)$, the overall effect of the skill will steer the swarm either towards the negative or positive direction; this generates the formation of two distinct clusters. Each cluster has an individual barycentric velocity: $\gamma = \frac{\gamma_1}{N} = \frac{\pm\beta}{N}$; this guarantees that the average barycenter remains unchanged.

Specifically, whenever the skills drift assumes the value $-\beta$, the skill has an overall propensity to stay at the rear of the swarm. Accordingly, all regular agents will, with high probability, be endowed with the rank-based drift $-g$. Hence the regular agents possess a clear tendency to be driven toward the negative direction on \mathbb{R} . Conversely, in presence of the $+\beta$ realization, the skill is very likely to belong to the group of leaders. This imposes to the remaining $(N-1)$ regular fellows to equally share, with alternations, the rank-based drift of $(N-1)g$ which drives the laggard. Therefore, with the $+\beta$ realization, the whole population is driven towards the positive direction of \mathbb{R} . We now can isolate three regimes depending on the strength β .

a) $\beta < \beta_c := \frac{N}{N-1}g$. In this regime, the skill is able to steer the whole population in one of the two possible directions while remaining itself attached to the swarm. In other words, a stationary probability measure exists for the distance between the agents (i.e. all agents evolve in a single flock). Note however that the presence of the skill breaks an initially uni-modal spatial repartition into a **bi-modal repartition**.

For three agents (i.e. one skill and two regular fellows), the resulting spatial repartition obtained by simulation is shown in Figure (1). In Figures (1) and (2), the skills position is represented in black, the others being displayed in grey.




Figure 1: Probability of spatial repartition of the agents at $t_{end} = 10$, numerical computation over 10^3 runs, with $N = 3$, $g = 1$, $\beta = 1.1 < \beta_c = \frac{3}{2}$.

b) $\beta_c \leq \beta < g \cdot N$. Here the strength of the ballistic noise precludes to reach a global stationary state for the inter-distance processes between the N agents. Indeed, the renormalized drift associated with the $+\beta$ realization of $dZ(t)$ (i.e. $\hat{\gamma}_1 = \gamma_1 + \beta$), violates the constraint Eq.(13). However, for the $-\beta$ realization, the constraint for $\hat{\gamma}_1 = \gamma_1 - \beta$ is fulfilled, implying that the skill remains flocked with its $(N-1)$ remaining fellows. An experiment with β chosen in this range, shows the spatial repartition in Figure (2) (left).




Figure 2: Left: End position distribution at $t_{end} = 10$, with $N = 3$, $g = 1$, $\beta = 2 > \beta_c$.

Right panel: Probability density of the position at $t_{end} = 10$ for $N = 3$, $g = 1$, $\beta = 4 > N \cdot g$. Both numerical computations include 10^3 runs.

c) Finally, for $\beta > g \cdot N$, the skill becomes highly turbulent and the tightness constraints are never fulfilled. The skill escapes from the flock and the resulting spatial repartition is typically shown in Figure (2) (right).

For all choices of the ballistic strength β , the numerical results intimately match the analytical predictions. In particular, the shill escapes from the flock when the critical ballistic strength $\beta = \beta_c$ is reached (for the $+\beta$ realization, or when $\beta = N \cdot g$ for the $-\beta$ realization of $dZ(t)$). In these non-stationary regimes, the shill quits the $(N - 1)$ - regular tight swarm with velocity $\mathcal{V}(\beta) \propto \beta$.

3.2 Distance Between Successive Agents

Complementary to the tightness constraints, let us now briefly discuss the stationary probability measure which characterizes the distances between successive agents. To this aim, we shall choose β to ensure the existence of a global stationary regime (i.e. ensures tightness), namely $0 \leq \beta < \beta_c = \frac{N}{N-1}g$.

Note however that using the results derived in (Ichiba T. and Fernhold, 2011), heterogeneous diffusion constants can also be analytically discussed. The distances separating consecutive agents defined in Eq.(6) can be explicitly computed by using the results summarized in section (2). For agent $Y_1(t)$ and in view of the extra β -drift induced by the shill ballistic driving, we are now led to define a couple of name-based drifts vectors as:

$$\gamma^\pm = \pm \left(\frac{N-1}{N}\beta \quad -\frac{\beta}{N} \quad \dots \quad -\frac{\beta}{N} \right). \quad (16)$$

The fulfillment of the tightness constraints given in Eq.(4) ensures that the $\lambda_{p,k}$ from Eq. (8) are strictly positive ($\forall p,k$). Hence, it results a couple of stationary probability densities (one for each realization $+\beta$ and $-\beta$), characterizing the inter-distance process $\Xi(t)$ (see Eq.(6)):

$$\begin{aligned} \mathbb{P}\{z \leq \Xi(t) \leq z + dz \mid \pm\beta \text{ is realized}\} &= \mathbb{P}_{\pm\beta}(z) dz \\ &= \left(\sum_{p \in \Sigma_N} \left(\prod_{k=1}^{N-1} \lambda_{p,k}^\pm \right)^{-1} \right)^{-1} \sum_{p \in \Sigma_N} \exp(-\langle \lambda_p^\pm, z \rangle) dz. \end{aligned} \quad (17)$$

The complete stationary probability density then reads:

$$\mathbb{P}(z) = \frac{1}{2} [\mathbb{P}_{+\beta}(z) + \mathbb{P}_{-\beta}(z)]. \quad (18)$$

4 USING A SHILL TO GUIDE THE COLLECTIVE CROSSING OF OBSTACLES

The shill $Y_1(t)$ of section (3) has so far been viewed as a mere trouble maker. However, for specific tasks to be achieved, a shills presence might become truly beneficial. Indeed, the presence of a shill can **softly control a swarm towards a preassigned target**.

To concretely illustrate this alternative view point, let us, once again, consider a collection of $(N - 1)$ identical drifted Brownian (regular) agents and one shill, say $Y_1(t)$, all diffusing on \mathbb{R} . From now on, we shall represent the evolution on \mathbb{R}^2 with the time being identified with the x -axis, and positions with the y -axis. At time $t = 0$, all N agents start at location $y_i(0) = 0$, $1 \leq i \leq N$. We may think of agents running on the x -axis with constant unit speed, as we therefore identify the x -axis with time. At the x -location T (see Figure 3), we introduce a solid wall (i.e. the obstacle) in which two-slots are drilled. The global objective for the N agents is to try to avoid the fatal collisions with the wall by traveling through one of the couple of slots. The two slots have width W and are symmetrically drilled at the ordinates $\pm D$, see Figure (3) for a sketch of the configuration.




Figure 3: Initial configuration for the soft control problem. The agents start their diffusion at $(t_0, y_0) = (0, 0)$.

In absence of shill, one has an homogeneous swarm (all agents are dynamically identical) and the N -swarm proceeds according to the rule defined in Eq.(1). As a consequence, the wall will be hit with high probability as, at time T , the probability density of the positions will be given by a collection of N centered Gaussians (we basically have N Brownian motions with constant drifts having their mean barycenter located at zero). As seen in section (3), the presence of a super-diffusive shill $Y_1(t)$ may strongly modify this nominal $\beta = 0$ picture. Indeed $Y_1(t)$, with suitable parameters β and g is able to steer the whole swarm with high preference to one of the slots. This basic and simple idea can be made fully quantitative as discussed below. In our simulations, we set $\beta = 0$ once the slots are crossed. Therefore, the global swarm continues its nominal path with a tight configuration after the obstacle.

4.1 Optimal Barycentric Driving for Suitable Choice of β

First, we select the $Y_1(t)$ parameters to ensure global tightness. This is achieved by defining $\beta < \beta_c = \frac{N}{N-1}g$. For a single realization of the $Z(t)$ noise, we already know that the average barycentric y -speed of the swarm is $\frac{\pm\beta}{N}$. Hence, at time T , the group barycen-

ter reaches the y -position located at $\frac{\pm\beta T}{N}$. The center of the slots being located at $\pm(D + \frac{W}{2})$, we therefore naturally require:

$$\frac{\pm\beta T}{N} = \pm \left(D + \frac{W}{2} \right) \Leftrightarrow \beta_{opt} := \frac{(D + \frac{W}{2}) \cdot N}{T}. \quad (19)$$

The choice β_{opt} , does not yet ensure that the skill $Y_1(t)$ itself remains attached to the flock. This second requirement can be achieved provided one has:

$$\begin{aligned} \frac{(D + \frac{W}{2}) \cdot N}{T} = \beta_{opt} < \beta_c = \frac{N}{N-1}g \\ \Leftrightarrow g > \frac{(D + \frac{W}{2}) \cdot (N-1)}{T} =: g_c. \end{aligned} \quad (20)$$

This choice of the couple parameters β_{opt} and $g > g_c$ now jointly ensures that i) the barycenter is steered towards one of the slots centers and ii) $Y_1(t)$ remains tightly attached to the swarm. The overall swarm's y -dispersion is itself dependent on the diffusion constants σ_i , which were here taken as ($\sigma_i := \sigma \forall i$).




Figure 4: Left panel: $N = 10$ agents, $W = 20$ and $D = 10$. Right panel: $N = 20$ agents, $W = 20$ and $D = 50$.

4.2 Adjusting the Spatial Dispersion Resolution for a Given Drift g

When the drift g is fixed, the ballistic component β cannot always be chosen to simultaneously ensure tightness and the collective drive into one of the slots. Keeping the barycentric driving $\beta = \beta_{opt}$, as defined in Eq.(19), three different scenarios are now possible:

a) $\beta = \beta_{opt} < \beta_c$. In this case, the skill nicely steers the group towards one of the slots while staying attached to the flock, as we already established in section 4.1.

b) $\beta = \beta_{opt} \geq \beta_c$. Here, $Y_1(t)$ escapes from the $(N-1)$ -flock with a constant drifting velocity. Two sub-cases have to be distinguished, i) drift $\beta - g$ resulting when $+\beta$ is realized by $dZ(t)$ and conversely ii) drift $(N-1)g - \beta$ for the alternative $-\beta$ case. In both cases, the regular agents evolve with an average drift γ_{ave} which reads:

$$\gamma = \begin{cases} \frac{\beta}{N} = \frac{(\beta-g) \cdot 1 + (N-1) \cdot \gamma_{ave}}{N} & \text{for } +\beta \\ -\frac{\beta}{N} = \frac{((N-1)g - \beta) + (N-1)\gamma_{ave}}{N} & \text{for } -\beta \end{cases}$$

$$\Leftrightarrow \gamma_{ave} = \begin{cases} \frac{g}{N-1} \\ -g \end{cases}. \quad (21)$$

c) $\beta = \beta_{opt} \geq \beta_c < N \cdot g$. Here, the skill remains attached to the flock for the $-\beta$ realization of $dZ(t)$, but escapes otherwise. This then leads to a mix of cases a) and b) depending on whether $+\beta$ or $-\beta$ is realized.

The computation of γ_{ave} from case b) shows that with g fixed such that $\beta_{opt} \geq \beta_c$, the skill escapes from the flock (hence, no stationary probability measure exists). The remaining $(N-1)$ regular fellows evolve with average speed γ_{ave} (which is β -independent). Whatever the values taken by β , the skill is never able to drive the swarm through one of the slots, the swarm's speed being only **g -dependent** if the skill is not attached to the swarm.

So far, only the swarm's directions has been considered. Obviously, the dispersion is also a determinant feature for efficient slots crossings. Here, not only g but the ratio $\frac{g}{\sigma}$ will be determinant. Clearly for small g values, the swarm dispersion will exhibit a clear tendency to exceed the slots widths, altering therefore the overall efficiency.

4.3 Multi-slots Configurations

So far, we did consider the capability of a skill to steer the swarm through a couple of slots. For multi-slots configurations, the skill construction used before naturally suggests to define more general skills to steer swarms in many different directions. This is achievable by replacing the ballistic noise driving the skill with more complex stochastic processes. Doing so however, the skill dynamics cannot be anymore represented by a simple diffusive stochastic differential equation like Eq.(11). As an illustration, consider a three symmetrical slots configuration for which one is naturally driven to introduce the following noise source:

$$dZ(t) = \begin{cases} \beta \tanh(\beta Z(t)) dt + dW(t) & \text{with prob. } \frac{2}{3}, \\ dW(t) & \text{with prob. } \frac{1}{3}. \end{cases} \quad (22)$$

The skill dynamics is taken as before, namely with probability $\frac{2}{3}$ it steers the swarm towards the positive or negative slots with ballistic parameter β and with probability $\frac{1}{3}$, it behaves as a regular agent driving the swarm on a centered path, see Figure (5).

For arbitrary number of slots and configurations, one can generalize our construction by suitably adjusting the properties of the noise source driving the skill.




Figure 5: Trajectories for β and g optimal, with $N_d = 3$ doors, under different conditions for the position/width of the doors.

5 CONCLUSIONS

It is a true challenge, to analytically discuss the swarm dynamics of heterogeneous interacting agents. Due to heterogeneity, ordinary analytical tools like the mean-field approach are to be ruled out a priori. Hence besides simulation experiments, very little hope remains for rigorous theoretical results. We think particularly to modeling approaches relying on statistical mechanics and phase transitions to explain the emergence of self-organized spatio-temporal patterns (i.e. *flocking*). However as our paper intends to show, facing inhomogeneous swarms problems, complete hope for analytical results should not be abandoned. The theoretical analysis, when achievable, definitely offers sources of inspiration for new and unexpected research axis.

In our present contribution, we barely scratched the wealth of analytical possibilities. Indeed, swarm heterogeneity has numerous origins, affecting the individual drifts and/or the variances, modeling the sensitivity of each agent to the external random environment. In parallel, heterogeneity can either be systematic, thus implying that each agent behaves differently, or can be limited to one or only a few individuals who exhibit singular behaviors. In the latter configuration, the emergent swarm dynamics can be affected, sometimes even strongly, by the exotic behavior of this (or these) individual(s). The influence of the exotic fellow(s) can hence be viewed as a *soft control* mechanism, either harmful or beneficial. The exotic insiders, acting as leaders (or as *shills* in economy) are not detected to be singular by the other fellows, offering the (politically frightening possibility!) to drive large swarms towards global goals known only to the manipulators. A formal analytical approach (complemented with simulations) to this general problematic is a truly fascinating challenge.

Further works include analysis of multiple shills influence, leading to the separation of the initial swarm into multiple flocks, one shill soft-controlling each flock. Generalization in two or three dimensions would also provide more realistic applications.

REFERENCES

- Banner A. D., R. F. and Karatzas, I. (2005). Atlas models of equity markets. *Ann. Appl. Prob.* 15, 2296-2330.
- Bialek W., Cavagna A., G. I. M. T. S. E. M. V. and Walczak, A. M. (2012). Statistical mechanics for natural flocks of birds. *PNAS* 13, vol. 109, 4786-4791.
- Chatterjee, S. and Pal, S. (2010). A phase transition behaviour for brownian motions interacting through their ranks. *Prob. Theory Relat. Fields* 147, 123-159.
- Dosseti, V. (2012). Cohesive motion in one-dimensional flocking. *Physica A: Math. Theor.* 45.
- Faria J. J., Dyer J. R. G., . . . and Krause, J. (2010). A novel method for investigating the collective behaviour of fish: Introducing 'robofish'. *Behavioral Ecology and Sociobiology* 8, vol. 64, 1211-1218.
- Gribovskiy A., Halloy J., D. J. L. H. B. and Mondada, F. (2010). Towards mixed societies of chickens and robots. *IEEE/RSJ 2010 International Conference on Intelligent Robots and Systems, Conference Proceedings* pp. 4722-4728.
- Han, J. and Wang, L. (2010). New strategy of the shill: Consistent moving. *Proceedings of the 29th Chinese Control Conference*, pp. 4459-4464.
- Hongler M. O., Filliger R., P. B. and Rodriguez, J. (2012). "Contemporary Topics in Mathematics and Statistics with Applications" in "On Stochastic Processes Driven By Ballistic Noise Sources". Ed. M. R. Adhikari.
- Hongler M. O., R. F. and Blanchard, P. (2006). Soluble models for dynamics driven by a super-diffusive noise. *Physica A* 370, 301-355.
- Ichiba T., Papathanakos V., B. A. I. K. and Fernhold, R. (2011). Hybrid atlas model. *Ann. Appl. Prob.* 21(2) 609-644.
- Pal, S. and Pitman, J. (2008). One dimensional brownian particles with rank-dependent drifts. *Ann. Appl. Prob.* 18 2179-2207.
- Rogers, L. C. G. and Pitman, J. W. (1981). Markov functions. *Ann. of Prob.* 9(4), 573-582.
- Vaughan R., Sumpter N., H. J. A. F. and Cameron, S. (2000). Experiments in automatic flock control. *Robotics and Autonomous Systems* 1, vol. 31, 109-117.
- Wang, L. and Guo, L. (2008). Robust consensus and soft control of multi-agent systems with noises. *Journal of Systems Science and Complexity* 3, vol. 21, 406-415.
- Wang X., J. H. and Han, H. (2011). Special agents can promote cooperation in the population. *PLoS ONE* 12, vol. 6.
- Wang Y., X.-Z. M. and Liu, J.-P. (2009). Soft control for swarm systems with simple attraction and repulsion functions. *Proceedings of the 2nd International Conference on Intelligent Networks and Intelligent Systems*, pp. 482-485.
- Yates C. A., Erban R., . . . and Sumpter, D. J. T. (2009). Inherent noise can facilitate coherence in collective swarm motion. *PNAS* 14, vol. 106, 5464-5469.